

SYSTEMATIC REVIEW

OPEN

Effect of Inhaled Volatile and IV Anesthetics on Biological Markers of Inflammation in Adult ICU and Thoracic Surgical Patients: A Systematic Review and Meta-Analysis

OBJECTIVES: Inhaled anesthetics may reduce alveolar and systemic inflammation in surgical and critically ill patients. This study aimed to perform a systematic review and meta-analysis comparing the effect of inhaled volatile and IV anesthetics on alveolar and plasma cytokines in patients with surgical or medical acute lung injury.

DATA SOURCES: Medline, Embase, and Cochrane CENTRAL databases from 2000 to July 2021.

STUDY SELECTION: Randomized control trials, prospective, and retrospective observational studies comparing inhaled volatile to IV anesthetics in ventilated adult patients with acute lung injury from lung resection or critical illness.

DATA EXTRACTION: A systematic review and meta-analysis was performed. Primary outcome was alveolar inflammatory cytokines levels that were meta-analyzed using a random effects model. Secondary outcomes were plasma inflammatory cytokine levels, mortality, pulmonary complications, and duration of hospital and ICU stay. The quality of studies was assessed using the Cochrane Risk of Bias tool for randomized control trials and the Cochrane Risk Of Bias In Non-randomized Studies of Interventions tool for retrospective cohort studies.

DATA SYNTHESIS: From 2522 screened studies, 28 (27 thoracic surgery and 1 ICU, $n = 4175$) were included. Meta-analysis of patients undergoing lung resection demonstrated lower levels of alveolar tumor necrosis factor- α (TNF- α) (standard mean difference 1.04; 95% CI, 0.32–1.77; $p < 0.01$; I^2 82%) and interleukin (IL)-6 (0.64; 95% CI, 0.52–0.75; I^2 0%; $p < 0.01$) at 1–2 hours in the inhaled anesthesia group, with no difference in other cytokines at various time points. The single ICU study demonstrated lower plasma TNF- α and IL-6 and alveolar TNF- α , IL-6, and IL-8 at 48 hours in patients sedated with sevoflurane compared with midazolam. Clinical outcomes were infrequently reported.

CONCLUSIONS: Limited evidence suggests that inhaled anesthesia may reduce proinflammatory cytokines TNF- α and IL-6 during lung resection and critical illness. Further studies are needed to clarify its effects on biological markers and clinical outcomes.

KEYWORDS: critical care; cytokine; inflammation; lung injury; lung resection; one-lung ventilation; thoracic surgery; volatile anesthetics

Inhaled volatile anesthetics are effective hypnotic agents that are capable of providing both surgical levels of anesthesia and sedation for critically ill patients (1, 2). They also possess excellent bronchodilator and anticonvulsant properties that are used to manage critically ill patients with medical

Soroush Rouhani, MD, FRCPC^{1,2}Sanchit Gupta, MD³Hira Raheel, MD^{1,2}Aggie Duan Gao, MD, FRCPC^{1,2}Ciara Hanley, MB, BCh, BAO,
FCAI, FJFICM^{1,4}Xingshan Cao, PhD⁵Alla Iansavitchene, MLIS⁶Brian H. Cuthbertson, MB, BCh,
MD, FRCA^{3,5,7,8}Marat Slessarev, MD, PhD,
FRCPC^{1,2}Ewan C. Goligher, MD, PhD,
FRCPC^{9,10,11,12}Aleksandra Leligdowicz, MD,
PhD, FRCPC^{1,2,13}Douglas D. Fraser, MD, PhD,
FRCPC^{2,14}Beverley A. Orser, MD, PhD,
FRCPC³Angela Jerath, MSc, FRCPC^{3,5,8,15}on behalf of the SAVE-ICU
Investigators

Copyright © 2025 The Authors. Published by Wolters Kluwer Health, Inc. on behalf of the Society of Critical Care Medicine. This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

DOI: 10.1097/CCE.0000000000001280



KEY POINTS

Question: The goal of this study was to perform a systematic review and meta-analysis comparing the effect of inhaled volatile and IV anesthetics on alveolar and plasma cytokines in patients with surgical or medical acute lung injury.

Findings: Meta-analysis of patients undergoing lung resection (27 studies) demonstrated significantly lower levels of alveolar tumor necrosis factor-alpha (TNF- α) and interleukin (IL)-6 at 1–2 hours in the inhaled anesthesia group compared with IV anesthetics. A single ICU study demonstrated lower plasma TNF- α and IL-6 and alveolar TNF- α , IL-6, and IL-8 at 48 hours in patients sedated with sevoflurane compared with midazolam.

Meaning: Limited evidence suggests that inhaled anesthesia may reduce proinflammatory cytokines TNF- α and IL-6 during lung resection and critical illness.

refractory status asthmatics and status epilepticus (3). Beyond their hypnotic properties, volatile anesthetics modulate immune function by suppressing proinflammatory cytokines such as tumor necrosis factor-alpha (TNF- α) and interleukin (IL)-6, and inhibiting toll-like receptor signaling (4). These effects are mediated through interactions with receptors expressed on neutrophils, macrophages, and lymphocytes, including gamma-aminobutyric acid type A, glycine, nicotinic acetylcholine, *N*-methyl-D-aspartate, and β 2 integrins (5). Although largely derived from preclinical and perioperative studies, this mechanistic evidence supports their potential immunomodulatory role in critical illness (6, 7).

In patients with lung injury from either surgical lung resection with one-lung ventilation (OLV) or acute respiratory distress syndrome (ARDS) requiring ICU admission, inhalational anesthetic agents may offer patient and health system benefits such as improved oxygenation, lower morbidity and mortality, and shorter duration of mechanical ventilation and ICU stay (6, 8–10). Given the distinct clinical contexts and pathophysiology of OLV and ARDS, we analyze them separately in this review. A prior meta-analysis of cardiac surgery patients showed that inhalational

anesthesia was associated with a lower risk of death, and pulmonary and non-pulmonary complications compared with IV anesthesia (9). These outcomes may in part be attributed to the immunomodulatory effects of inhalational anesthetics, including the reduction of proinflammatory cytokines such as TNF- α , IL-6, and IL-8. We selected these cytokines as primary markers of interest because they are central to the pathogenesis of acute lung injury—driving neutrophil recruitment, vascular permeability, and systemic inflammation—and are among the most consistently reported biomarkers in studies of inhaled anesthetics (4). Inhalational agents also exert broader effects on immune cells, including macrophages, lymphocytes, and neutrophils, through interaction with cellular receptors that modulate inflammatory signaling pathways (4).

Although several studies have examined the effect of inhalational compared with IV anesthesia and sedation on inflammatory cytokines, these data have not been systematically summarized across studies. This systematic review and meta-analysis evaluate the effect of inhaled (intervention) vs. IV (comparison) anesthetics on levels of TNF- α , IL-6, and IL-8 (outcomes), in adult patients with acute lung injury (population) resulting from either surgical OLV or ARDS.

METHODOLOGY

This protocol was developed using the Preferred Reporting Items for Systematic Review and Meta-Analysis Protocols (11) and registered with PROSPERO (CRD42023444105).

Study Eligibility Criteria

Inclusion Criteria. (1) Studies in mechanically ventilated adult patients 18 years old or older admitted to critical care settings with acute lung injury/ARDS, and surgical patients undergoing OLV, (2) randomized controlled trials, (3) prospective and retrospective observational studies, (4) studies that compare inhaled volatile and IV sedative agents (studies with only one group were not included), (5) inhalational agents include sevoflurane, isoflurane, desflurane, enflurane, and (6) published in English only from 2000 to July 2021. We limited our search to this date range to ensure applicability to contemporary clinical practice, as anesthetic agents and perioperative management strategies have evolved significantly over the past 2

decades, with a shift from older agents such as halothane to modern volatile anesthetics like isoflurane, sevoflurane, and desflurane.

Exclusion Criteria. (1) Quasi-randomized studies, crossover studies, case reports, and case series, (2) preclinical studies and animal studies, (3) studies involving noninvasive ventilation, (4) studies involving pediatric or pregnant patients, (5) abstracts only, and (6) editorial articles without original data.

Outcomes

The primary outcomes were cytokine levels measured in either bronchoalveolar lavage (BAL) fluid or plasma. Although BAL may provide a more direct assessment of local pulmonary inflammation, plasma cytokine levels are more routinely measured and clinically relevant; therefore, we report BAL and plasma data separately. Specific cytokines studied included proinflammatory cytokines TNF- α , IL-6 and IL-8, and anti-inflammatory cytokine IL-10. We selected TNF- α , IL-6, IL-8, and IL-10 as cytokines of interest based on their central role in the inflammatory response associated with acute lung injury, with TNF- α , IL-6, and IL-8 representing key proinflammatory mediators and IL-10 serving as a principal anti-inflammatory regulator. These cytokines are also the most consistently reported across relevant studies, facilitating data comparison and synthesis. In addition to these outcomes, reviewers extracted study details (year, location, number of patients, funding), patient demographics (age, sex), population (thoracic surgery or acute lung injury/ARDS), ventilation details, anesthetic agent use/dose, duration of anesthesia, all cytokines measured and time points for measurements (when available). Secondary outcomes included plasma levels of cytokines TNF- α , IL-6, IL-8, and IL-10, as well as clinical outcomes such as duration of ventilation, ventilator-free days, pulmonary complications (e.g., bronchospasm, atelectasis, pneumonia, pulmonary edema, ARDS), length of ICU and hospital stay, and death reported at any time points. We aimed to quantify clinical outcomes from these studies but this was not feasible given the paucity and heterogeneity of clinical outcome information after data extraction. Primary authors were not contacted for missing information. Quality assessment of all included studies was performed. The Cochrane Risk of Bias (RoB) tool was used for randomized control trials. Retrospective cohort studies were assessed using the Cochrane Risk

Of Bias In Non-randomized Studies—of Interventions (ROBINS-I) tool.

Electronic Search

Systematic literature searches were conducted in MEDLINE, Cochrane Central Register of Controlled Trials, and Embase (all via Ovid platform) electronic databases from 2000 to July 2021 and were restricted to adults (older than 18 years). The search was conducted by a medical information specialist with formal training in search strategies for systematic reviews in consultation with the review authors. Only English studies were reviewed. Studies from animal experiments were excluded. Review articles were cross-referenced for additional studies. Authors were not contacted for unpublished data. A sensitive search strategy was developed by combining synonymous searches comprised of controlled vocabularies, such as Medical Subject Headings in Medline or EMTree descriptors in Embase, and free-text terms into the search blocks of two distinct permutations: 1) Volatile/Inhalation anesthetics + (Critical/Intensive care or Thoracic Surgery requiring intubation and ventilation or One-Lung Ventilation) + IV Anesthesia + Inflammation, and 2) Volatile anesthetics + ventilation + IV agents. The search strategies were modified for each database using various types of thesaurus terms and fields within the database. To identify additional relevant studies that met inclusion criteria, the reference lists of the relevant studies were examined, and “clinicaltrials.gov” (National Institutes of Health) and the World Health Organization International Clinical Trials Registry Platform were searched to identify unpublished trials.

Study Selection and Extraction

Seven reviewers (S.R., S.G., C.H., H.R., A.G., A.J., M.S.) independently screened abstracts for study eligibility criteria. All studies were reviewed by two independent individuals and conflicts were resolved by a third reviewer. Full-text editions were obtained for all selected studies. Data extraction was performed using a standardized extraction tool in Excel and confirmed by a second independent reviewer. A plot digitizing tool (WebPlotDigitizer, San Francisco, CA) was used to extract quantitative data from graphical displays when numerical data were unavailable (12). Complete data will be made available upon reasonable request to the authors.

Statistical Analysis

Given the distinct clinical contexts and pathophysiology of OLV and ARDS, we analyzed these two patient cohorts separately. Descriptive statistics were used to describe studies using mean (SD) and frequency (percentage) for continuous and categorical variables respectively. Cytokines measured for patients undergoing thoracic surgery or admitted to ICU with acute lung injury/ARDS were separately described. Given limited data for ICU patients, a meta-analysis was performed only for patients undergoing thoracic surgery. The inflammatory cytokines TNF- α , IL-6, IL-8, and IL-10 were analyzed separately with levels measured at baseline (before incision), and then at 1–2 hours and 3–4 hours; these time points were chosen after reviewing common timing of cytokine level measurements, and time points of 1–2 hours and 3–4 hours coincided with performing OLV and two-lung ventilation during routine lung/esophageal resection. All levels of cytokines were compared between the IV and inhaled groups using mean difference (95% CI) in a meta-analysis. The meta-analysis used a random effects model with restricted maximum likelihood to estimate between study variance, and CIs calculated using Knapp-Hartung adjustment. Heterogeneity was assessed using I^2 (moderate heterogeneity I^2 25–50%, severe heterogeneity I^2 > 75%) and Cochran's Q test, and invariance methods to assess study weighting. If the full-text article had data presented as median, interquartile range, and range, the mean and SD were calculated using the methods outlined by Wan et al and Cochrane Handbook (13, 14). Meta-analyses were conducted using R studio, Version 2023 12.1 (R Foundation for Statistical Computing, Vienna, Austria). The number of available studies reporting cytokine levels at the above time points differed between the cytokines with a maximum of eight studies. Assessment of publication bias via funnel plots and Egger's statistic required a minimum of ten studies using our statistical software, and was thus not performed; however, we acknowledge the potential for such bias as a limitation of our findings.

RESULTS

Study Cohort

The search identified 2522 studies and full-text assessment was performed on 433 studies. Of these, 28

studies met eligibility with 27 thoracic surgery ($n = 4,125$) and one ICU ARDS ($n = 50$) study (**Fig. 1**) (6, 8, 15–40). All studies were single center with patient numbers ranging from 20 to 555 and compared sevoflurane to propofol in 86% surgical studies, and sevoflurane to midazolam in the single ICU acute lung injury study (**Supplemental Table 1**, <https://links.lww.com/CCX/B524>). Study patient characteristics included 56% males with a mean (SD) age of 54.1 (11.8) in the thoracic surgery group, and 72% males with a mean age of 64.5 (12) in the ICU patients. Mean duration of drug use was 4.7 (1.6) hours for thoracic surgical patients and 48 hours for the ICU patients (**Supplemental Table 1**, <https://links.lww.com/CCX/B524>).

Thoracic Surgery Patients

Serial change in cytokines TNF- α , IL-6, IL-8, and IL-10 between IV and inhaled anesthetic groups are shown in **Figure 2**.

The number of studies measuring TNF- α included in our meta-analysis were 6 ($n = 390$) (6, 16, 19, 25, 29, 30) and 6 ($n = 514$) (6, 15, 19, 21, 25, 30) in BAL and plasma samples respectively. At baseline, there was no significant difference in BAL and plasma TNF- α levels. At 1–2 hours, TNF- α levels were higher in the IV group in Five of six BAL studies ($n = 390$) with the meta-analysis showing a significant mean difference of 1.04 (95% CI, 0.32–1.77; $p < 0.01$; I^2 82%) (6, 16, 19, 25, 29, 30). Plasma TNF- α levels were higher in the IV group in Three of six studies ($n = 514$) (6, 15, 19, 21, 25, 30) with no significant difference overall (**Table 1**). At the 3–4 hours timepoint, one study from BAL samples (total 2 studies, 80 patients) (25, 29) and two studies among plasma samples (total 3 studies, 258 patients) (15, 21, 25) demonstrated higher TNF- α levels in the IV group compared with the inhaled group but the pooled meta-analysis revealed no significant difference overall (**Table 1**).

The number of studies measuring IL-6 included in our meta-analysis were 5 ($n = 350$) (6, 16, 19, 30, 32) and 9 ($n = 630$) (6, 15, 19, 21, 23, 27, 30, 33, 36) in BAL and plasma samples, respectively. At baseline, there was no significant difference in BAL or plasma IL-6 between groups. At 1–2 hours, BAL IL-6 levels were higher among IV groups in three of five studies ($n = 350$) (6, 16, 19, 30, 32) with a significantly higher




Figure 1. Preferred Reporting Items for Systematic Review and Meta-Analysis study flowchart.

pooled mean difference (0.64; 95% CI, 0.52–0.75; I^2 0%; $p < 0.01$). Plasma IL-6 levels were higher in the IV group in two of seven studies ($n = 562$) (6, 15, 19, 21, 27, 30, 33), whereas four studies showed no difference, and one study showed higher levels among the inhaled anesthetic arm. Overall the meta-analysis revealed no significant difference between the groups. At 3–4 hours, IL-6 was measured in BAL in only one study (36), and in plasma in five studies (15, 21, 23, 27, 36), with pooled meta-analysis data demonstrating no overall difference.

29, 36) and three studies with plasma samples (25, 27, 36), and the pooled meta-analysis revealed no significant difference overall.

The number of studies measuring anti-inflammatory cytokine IL-10 that were included in the meta-analysis were 3 ($n = 236$) (6, 30, 36) and 5 ($n = 350$) (6, 17, 21, 30, 33) in BAL and plasma samples, respectively. At 1–2 hours, one of two studies ($n = 216$) (6, 30) measuring BAL levels showed higher levels of IL-10 in the inhaled group but the pooled mean difference was not significantly different between groups.

The number of studies measuring IL-8 included in the meta-analysis were 6 ($n = 390$) (6, 16, 25, 29, 30, 32) and 4 ($n = 286$) (6, 19, 30, 36) in BAL and plasma samples, respectively. At baseline, there was no significant difference in BAL or plasma IL-8 between groups. At 1–2 hours, IL-8 levels were higher among the IV anesthetic group in three of six BAL studies ($n = 278$) (6, 16, 25, 29, 30, 32); however overall, there was a non-significant standard mean difference of 1.50 (95% CI, -0.27–3.26; $p = 0.08$; I^2 96.6%). Similarly, two of three studies (6, 25, 30) with plasma samples showed higher values of IL-8 among the IV anesthetic group compared with the inhaled anesthetic group with no significant pooled mean difference overall. IL-8 was measured at 3–4 hours in three studies with BAL samples (25,




Figure 2. Temporal changes in mean bronchoalveolar lavage (BAL) and plasma cytokine levels across studies. Units are in pg/mL. IL = interleukin, TNF- α = tumor necrosis factor-alpha.

Similarly, two of four studies (21, 30) showed higher plasma levels of IL-10 in the IV group at 1–2 hours, with no significant pooled mean difference between groups overall. Two studies measured BAL IL-10 levels ($n = 50$) (29, 36) and four studies measured plasma IL-10 levels ($n = 170$) (17, 21, 27, 33) at 3–4 hours. Pooled

stay were recorded in 6 ($n = 406$) (6, 16, 23, 25, 35, 38) and 7 ($n = 2,177$) (6, 23, 25, 26, 34, 38, 40) studies, respectively. ICU length of stay was significantly longer in the propofol arm of three studies (16, 25, 38), with the remainder showing no significant difference between groups. Among studies reporting hospital length

data showed no significant difference in BAL or plasma IL-10 levels between groups.

Several other biomarkers were presented infrequently among thoracic surgery studies, including inflammatory factor matrix metalloproteinase-9 (MMP-9) and metrics of oxidative stress such as malondialdehyde (MDA). MMP-9 was only reported in plasma samples in two studies (37, 39) demonstrating increased plasma concentration in the sevoflurane group at chest closure. MDA was measured in plasma samples in two studies (18) with only Yang et al (39) demonstrating significantly higher levels in the sevoflurane arm compared with propofol after chest closure. A comprehensive list of reported biomarkers for each study is presented in Supplemental Table 1 (<https://links.lww.com/CCX/B524>).

Clinical outcomes were infrequently reported among the thoracic surgery studies. Of the 27 studies, ICU and hospital length of

TABLE 1.

Summary of Heterogeneity and Standard Mean Difference of Bronchoalveolar Lavage and Plasma Cytokine Levels in Patients Undergoing IV vs. Inhaled Anesthesia

Biomarker	Timepoint (hr)	Studies/ Participants	IV vs. Inhaled Standard Mean Difference (95% CI)	<i>p</i>	<i>I</i> ² (%), Cochran Q, <i>p</i>)
Bronchoalveolar lavage data					
TNF- α	Baseline	5/336	0.23 (−0.43 to 0.88)	0.39	59.5, 9.87, 0.04
	1–2	6/390	1.04 (0.32 to 1.77)	0.01 ^a	82.27, 80, < 0.01
	3–4	2/80	0.79 (−7.16 to 8.74)	0.43	85, 6.66, 0.01
IL-6	Baseline	5/316	0.15 (−0.24 to 0.54)	0.33	36.2, 6.27, 0.18
	1–2	5/350	0.64 (0.52 to 0.75)	< 0.01 ^a	0, 0.58, 0.96
	3–4	1/20	−0.11 (−0.99 to 0.76)	0.8	Not available
IL-8	Baseline	6/356	0.22 (−0.07 to 0.52)	0.11	12.9, 5.74, 0.33
	1–2	6/390	1.50 (−0.27 to 3.26)	0.08	96.6, 146.21, < 0.01
	3–4	3/100	1.02 (−7.09 to 9.12)	0.64	97.1, 68.50, < 0.01
IL-10	Baseline	3/236	−0.27 (−0.88 to 0.34)	0.2	14.1, 2.33, 0.31
	1–2	2/216	−0.79 (−10.85 to 9.27)	0.5	95, 20.02 < 0.01
	3–4	2/50	0.48 (−3.40 to 4.35)	0.36	8.9, 1.10, 0.29
Plasma data					
TNF- α	Baseline	6/514	−0.11 (−0.45 to 0.23)	0.44	42.9, 8.76, 0.12
	1–2	6/514	0.67 (−0.42 to 1.76)	0.18	93.4, 76.16, < 0.01
	3–4	3/258	0.94 (−4.13 to 6.02)	0.51	96.7, 60.56, < 0.01
IL-6	Baseline	9/630	−0.16 (−0.55 to 0.22)	0.36	72.3, 28.85, < 0.01
	1–2	7/562	0.24 (−0.69 to 1.18)	0.55	94.9, 117.31, < 0.01
	3–4	5/312	0.44 (−1.08 to 1.97)	0.46	91.1, 45.19, < 0.01
IL-8	Baseline	4/286	−0.05 (−0.39 to 0.29)	0.66	0, 2.45, 0.48
	1–2	3/266	1.70 (−2.49 to 5.88)	0.22	95.1, 40.52, < 0.01
	3–4	3/106	0.77 (−2.03 to 3.58)	0.36	90.2, 20.32, < 0.01
IL-10	Baseline	5/350	0.31 (−1.75 to 2.38)	0.69	94.8, 76.97, < 0.01
	1–2	4/292	0.01 (−0.33 to 0.34)	0.95	0, 2.38, 0.50
	3–4	4/170	1.25 (−2.29 to 4.80)	0.34	96.0, 74.51, < 0.01

IL = interleukin, TNF- α = tumor necrosis factor-alpha.

^aStatistical significance ($p < 0.05$).

Data are presented as standard mean difference (95% CI).

of stay, there was no significant difference observed between groups. Ventilator-free days were not recorded in any of the included studies. Postoperative pulmonary complications were reported in 11 studies ($n = 2,200$) (6, 16, 23–27, 35, 36, 38, 40), with varying definitions and reporting standards. Among these, three studies (6, 16, 25) showed an increased rate of postoperative complications in the propofol group, with the remainder showing no difference between groups. Eight studies ($n = 2,620$) reported on death after surgery during hospitalization, at 30 days or at 1 year (6, 16, 23–25, 27, 38, 40). Only one study (6) reported significantly increased mortality in the IV group, with the remaining seven studies showing no significant difference in mortality between groups. Duration of ventilation was infrequently reported, and was estimated based on other reported metrics such as duration of anesthesia, or total surgical time, with no difference noted between groups (**Supplemental Table 2**, <https://links.lww.com/CCX/B524>).

ARDS Patients in the ICU

A single ARDS study ($n = 50$) (8) measured several biomarkers including s-RAGE (soluble receptor for advanced glycation end-products), TNF- α , IL-6, and IL-8. s-RAGE is released from type I alveolar cells in ARDS (41) and was shown to be significantly lower after 48 hours of sevoflurane therapy compared with IV midazolam ($p = 0.02$ and 0.04 in plasma and BAL, respectively). This study also showed significantly lower alveolar levels of TNF- α , IL-6 and IL-8, and plasma levels of TNF- α and IL-6 in the sevoflurane group compared with the midazolam group at 48 hours. Clinical outcomes measured in this study included pulmonary oxygenation, which was significantly higher at 48 hours among the sevoflurane group compared with midazolam ($\text{Pao}_2/\text{Fio}_2$ ratio 205 [56] vs. 166 [59], $p = 0.04$ for post hoc comparisons). Other clinical outcomes, such as ICU length of stay, ventilator-free days and death, were not significantly different between treatment arms.

Risk of Bias

The risk of bias assessment performed for each study is summarized in **Supplemental Table 3** (<https://links.lww.com/CCX/B524>). Of the 27 included studies, 25 randomized control trials (RCTs) were assessed by the

RoB tool, and two retrospective cohort studies were assessed by the ROBINS-I tool. Among the 25 RCTs, nine studies were identified as low risk of bias, and seven were of uncertain risk. The remainder showed intermediate to high risk, in the domains of deviation from intended interventions (lack of blinding in most RCTs), and bias in missing outcomes/measurement of outcomes. The two non-randomized studies were deemed low risk of bias overall.

DISCUSSION

This systematic review and meta-analysis evaluates and compares the levels of inflammatory cytokines in BAL and plasma between patients receiving inhaled vs. IV anesthetics in patients with acute lung injury resulting from either surgical OLV or ARDS. Our meta-analysis demonstrated higher levels of inflammatory cytokines TNF- α and IL-6 in BAL samples among surgical patients receiving IV compared with inhaled anesthesia—particularly at the 1–2 hours mark during the point of maximal inflammatory insult of OLV and lung resection. No significant differences were seen in pooled data for plasma concentrations of inflammatory cytokines. In a single study of ventilated ARDS patients, sevoflurane, when compared with midazolam reduced alveolar levels of inflammatory cytokines s-RAGE, TNF- α , IL-6 and IL-8, and plasma levels of TNF- α and IL-6.

Both proinflammatory (e.g., TNF- α , IL-6, IL-8) and anti-inflammatory cytokines (e.g., IL-10) are implicated in the pathogenesis of acute lung injury and ARDS (42–46). Local elevations in inflammatory cytokines are seen almost immediately after an initial insult; in the setting of lung injury from OLV or ARDS, this signal would first be expected in alveolar fluid (47–49). TNF- α is released early in the proinflammatory cascade and is known to amplify downstream cytokines, including IL-6, IL-8, and several others (44, 50). In certain lung pathologies (e.g., COVID-19), over-production of TNF- α is persistent and related to poorer prognosis (51). This study may support the hypothesis that inhaled anesthetics such as sevoflurane and isoflurane help attenuate acute alveolar inflammatory responses by modulating the expression of pro-inflammatory cytokines TNF- α and IL-6. Potential mechanisms implicated in this include activation of mitochondrial K-adenosine triphosphate channels, which preserves mitochondrial function and reduces

cellular injury during inflammatory stress. Inhaled anesthetics also inhibit toll-like receptor signaling, suppress NF- κ B activation, and modulate protein kinase pathways, providing a plausible biological basis for their ability to reduce cytokine release and mitigate inflammatory lung injury (4, 5, 52, 53).

There was no significant difference in alveolar or plasma levels of proinflammatory cytokine IL-8 and anti-inflammatory cytokine IL-10 in our meta-analysis.

The reduction in BAL concentrations of inflammatory cytokines with inhaled vs. IV anesthesia did not translate in to corresponding changes in plasma cytokine levels within our meta-analysis. A number of hypotheses are postulated. First, the inflammatory insult of OLV may be insufficient for inducing a systemic inflammatory response. Second, the relatively narrow observation time (up to 4 hr) may be insufficient to detect changes in systemic inflammatory response. Cytokine measurements at later time points such as 24 hours after surgery showed no difference between anesthetic groups but were reported by very few studies that precluded meaningful pooling of data (6, 21, 23, 33, 36). Only a single study gathered data beyond the 24 hours time point (26). Third, the paucity of studies measuring plasma cytokine concentrations in surgical patients undergoing OLV also limits the power to detect any such differences. Fourth, the effect of inhaled anesthetic concentration and duration of exposure may impact the systemic inflammatory response, but this was not studied. This is best illustrated in the single ARDS study, where longer exposure to inhaled anesthetic and longer observation period enabled detection of inflammatory response in both BAL and plasma samples, as well as detection of attenuating effect in inhaled anesthetics arm at 48 hours (8). These explanations remain hypotheses, and the current data are insufficient to directly evaluate them. Further studies with standardized sampling timepoints, larger sample sizes, and longer follow-up are needed to better understand compartment-specific cytokine dynamics and the systemic effects of inhaled anesthetics.

An important finding of this review is methodologic heterogeneity with respect to the type of markers being measured, assays that are used, and variable measurement time points. It also demonstrates a marked gap in reporting clinical correlation of biomarker findings with patient and health system outcomes such as mortality, morbidity and duration of ventilation and ICU

stay. Although the clinical implications remain uncertain, reduced BAL cytokine levels may reflect attenuated pulmonary inflammation, which could contribute to improved gas exchange, less lung injury, and potentially shorter durations of ventilation and ICU stay. Future studies should aim to correlate these biomarker changes with clinical outcomes.

Overall, there was paucity of studies reporting standardized and comprehensive datasets. Given the growing interest that inhaled anesthetics possess immunomodulatory properties, our study highlights the urgent need to mandate imbedding measurement of cytokines and other inflammatory biomarkers into future clinical trials and developing standards for collecting, measuring and reporting these biomarkers as well as related patient and health system outcomes. Future studies should also consider assessing more comprehensive biomarker panels such as proteomics or RNA sequencing.

Additional limitations of this study include the low number of available studies, small sample size, and lack of data on clinical and surgical complexity. We attempted to mitigate this by selecting the most commonly represented cytokines, and in summarizing the data in the most parsimonious way possible, particularly with respect to amalgamating common time points among studies. Due to the paucity of data, a meta-analysis of ICU ARDS patients was not possible. Given extensive heterogeneity in reporting of patient and health system outcomes, a meta-analysis of these variables was also not performed.

Despite these limitations, our study suggests the postulated anti-inflammatory properties of inhaled anesthetics as a possible plausible biologic mechanism for improved outcomes in patients with lung injury from thoracic surgery. Building on knowledge regarding TNF- α dynamics and its role in disease progression (44, 50), this study accordingly highlights a rationale for earlier administration of inhaled anesthetics to improve clinical and biochemical outcomes. Furthermore, it highlights the importance of imbedding measurement of biomarkers into clinical trials to provide mechanistic information in how clinical interventions such as type of anesthetic agents can impact biologic processes and alter the pathophysiology of underlying disease states. Importantly, this may inform the development of future targeted therapeutic interventions. Our study identifies a need for standardization of the methodologic aspects

of biomarker research in this space including choice of biomarker assays, timing of sample collections, and routine reporting of core patient and health system outcomes along with the biomarker data. Finally, it highlights the importance of continuing work in characterizing the immunomodulatory effects of inhaled anesthetics, as novel inflammatory biomarkers are discovered and integrated in clinical research.

This work supports several active large clinical trials (NCT05327296, NCT05312385, NCT04341350, NCT04415060) that will improve our knowledge on the effect of inhaled anesthetics on patient outcomes and help establish protocols for the delivery of inhaled anesthetics in the ICU, guide best practices for administration, ventilator optimization, and troubleshooting. To facilitate broader adoption of inhalational anesthetics, it is essential to collaborate with anesthesia critical care specialists and anesthesiologists to ensure safe and effective implementation (3). In addition, surveying ICU providers to identify barriers, logistical challenges, and educational needs will be crucial in developing tailored strategies that support the integration of inhaled anesthetics into routine critical care practice.

CONCLUSIONS

In patients undergoing OLV for thoracic surgery, inhaled anesthesia was associated with attenuation of proinflammatory cytokines TNF- α and IL-6 compared with IV anesthesia. In ICU patients with ARDS, one RCT reported a similar effect, although this finding remains preliminary and should be interpreted with caution given the limited evidence base. Given the paucity of data and significant heterogeneity among studies, it is not clear if the same effect extends to other reported cytokines, and if these effects translate into meaningful improvement inpatient or health system outcomes. Future research should strive toward standardizing collection and analysis of inflammatory biomarkers, linking biomarker data to standard patient (mortality, morbidity) and health system (duration of ventilation and ICU stay) outcomes, and use of novel inflammatory biomarker assays as they become available for clinical research.

ACKNOWLEDGMENTS

The SAVE-ICU Study Investigators are as follows: Sunnybrook Health Sciences Centre: Martin Chapman, Damon Scales, Julie Nardi, Beth Linseman, Phil Lau, Eily

Shaw, Lilia Kaustov; Centre hospitalier de l'Université de Montréal (CHUM), Montréal, Canada: Emmanuel Charbonney, Michaël Chassé, Martin Girard, Antoine Halwagi, Guillaume Plourde, Han Ting Wang, Francois Carrier; Centre hospitalier universitaire de Sherbrooke (CHUS): Frédérick D'Aragon, Maxime Tissot-Therrien, François Lamontagne, Dominique Bérard; Centre intégré universitaire de santé et de services sociaux du Nord-de-Ille-de-Montréal (Sacre Coeur): Yiorgos Alexandros Cavayas, Martin Albert, Francis Bernard, Karim Serri, David Williamson, Virginie Williams, Patricia Martinez Barrios, Sabrina Araujo De Franca; Institut Universitaire de Cardiologie et de Pneumologie de Québec, Québec City, Canada: Etienne J. Couture, Adreanne Côté, Mathieu Simon, Pierre-Alexis Lépine, Dave Gleeton; London Health Sciences Centre: John Basmaji, Marat Slessarev, Ahmed Hegazy, Tracey Bentall, Eileen Campbell; Michelle Stephens; Tim Winterburn; McGill University Health Centre: Roupén Hatzakorzian, Kosar Khwaja, Peter Goldberg, Jason Shahin, Jeremy Richard Grushka; The Ottawa Hospital: Jonathan Hooper, Sherissa Microys, Hilary Meggison, David Leilipovitz, Jessica Haines, Irene Watpool, Rebecca Porteous, Stephanie Dunster, Sylvie Bourbonnais; Toronto General Hospital: Ewan Goligher, Lorenzo Del Sorbo, Niall Ferguson, Philip Graham, Michael Long, Tina Yan, Denise Morris, Maria Kobylecky, Hesham Abdelhady, Tina Romagnuolo; Toronto Western Hospital: Ian Randall, Elizabeth Wilcox, Bourke Tillman; University of Alberta Hospital: Michael Jacka, Oleksa Rewa, Vincent Lau, Sean Bagshaw, Nadia Baig. Grey Nuns Community Hospital: Janek Senaratne, Lazar Milovanovic, Tammi Breymann, Larissa Johnson, Anushka Jayasekara, Bernadette Fernando, Paula Sharman, Vanessa Dolimount, Jennifer Hamilton, Mehvash Qureshi.

1 Department of Medicine, London Health Sciences Centre, London, ON, Canada.

2 Schulich School of Medicine and Dentistry, Western University, London, ON, Canada.

3 Department of Anesthesiology and Pain Management, Temerty Faculty of Medicine, University of Toronto, Toronto, ON, Canada.

4 Department of Anaesthesia and Critical Care, University Hospital Galway, Galway, Ireland.

5 Sunnybrook Research Institute, Sunnybrook Health Sciences Centre, Toronto, ON, Canada.

6 London Health Sciences Centre, Corporate Academics, London, ON, Canada.

- 7 Department of Critical Care Medicine, Sunnybrook Health Sciences Centre, Toronto, ON, Canada.
- 8 Institute of Health Policy, Management and Evaluation, University of Toronto, Toronto, ON, Canada.
- 9 Interdepartmental Division of Critical Care Medicine, University of Toronto, Toronto, ON, Canada.
- 10 Department of Medicine and Physiology, University of Toronto, Toronto, ON, Canada.
- 11 Critical Care Section, University Health Network, Toronto, ON, Canada.
- 12 Toronto General Hospital Research Institute, Toronto, ON, Canada.
- 13 Robarts Research Institute, Western University, London, ON, Canada.
- 14 Department of Pediatrics, London Health Sciences Centre, London, ON, Canada.
- 15 Institute for Clinical Evaluative Sciences, Toronto, ON, Canada.

The SAVE-ICU Study Investigators are listed in the Acknowledgments.

Supplemental digital content is available for this article. Direct URL citations appear in the printed text and are provided in the HTML and PDF versions of this article on the journal's website (<http://journals.lww.com/ccejjournal>).

Dr. Jerath was involved in study concept, design study protocol, data extraction, data analysis, data interpretation, drafting the first article, and critical revision of the article. Dr. Rouhani was involved in data extraction, data analysis, data interpretation, drafting the first article, and critical revision of article. Drs. Gupta, Raheel, and Gao were involved in data extraction and article revision. Dr. Hanley was involved in study concept, design study protocol, data extraction, and article revision. Dr. Cao was involved in data analysis, data interpretation, and article review. Ms. Iansavitchene was involved in design literature search strategy and revision of article. Dr. Leligdowicz was involved in article review and critical revision. Dr. Slessarev was involved in conception, design study protocol, data analysis, data interpretation, drafting first article, and critical revision of article. Drs. Goligher, Fraser, Leligdowicz, Orser, and Cuthbertson were involved in article review and critical revision.

Drs. Jerath and Cuthbertson are supported by the Merit Award, Department of Anesthesiology and Pain Medicine, University of Toronto. Drs. Jerath, Slessarev, and Cuthbertson have received grant funding from the Canadian Institute of Health Research and the Ontario Government. Drs. Jerath and Cuthbertson received funding from Alternate Funding Plan, Ministry of Ontario. Drs. Jerath, Slessarev, and Cuthbertson received funding Government of Ontario. Drs. Slessarev and Jerath have an ongoing clinical trial examining the effects of inhaled anesthetics on cognitive outcomes in critically ill adult patients. Dr. Leligdowicz is a recipient of the Wolfe-Western Fellowship award.

For information regarding this article, E-mail: srouh035@uottawa.ca

REFERENCES

1. Jerath A, Ferguson ND, Cuthbertson B: Inhalational volatile-based sedation for COVID-19 pneumonia and ARDS. *Intensive Care Med* 2020; 46:1563–1566
2. Jerath A, Ferguson ND, Steel A, et al: The use of volatile anesthetic agents for long-term critical care sedation (VALTS): Study protocol for a pilot randomized controlled trial. *Trials* 2015; 16:560
3. Gorsky K, Cuninghame S, Jayaraj K, et al: Inhaled volatiles for status asthmaticus, epilepsy, and difficult sedation in adult ICU and PICU: A systematic review. *Crit Care Explor* 2024; 6:e1050
4. Stollings LM, Jia LJ, Tang P, et al: Immune modulation by volatile anesthetics. *Anesthesiology* 2016; 125:399–411
5. Yuki K, Eckenhoff RG: Mechanisms of the immunological effects of volatile anesthetics: A review. *Anesth Analg* 2016; 123:326–335
6. de la Gala F, Piñeiro P, Reyes A, et al: Postoperative pulmonary complications, pulmonary and systemic inflammatory responses after lung resection surgery with prolonged one-lung ventilation. Randomized controlled trial comparing intravenous and inhalational anaesthesia. *Br J Anaesth* 2017; 119:655–663
7. O'Gara B, Talmor D: Lung protective properties of the volatile anesthetics. *Intensive Care Med* 2016; 42:1487–1489
8. Jabaudon M, Boucher P, Imhoff E, et al: Sevoflurane for sedation in acute respiratory distress syndrome. A randomized controlled pilot study. *Am J Respir Crit Care Med* 2017; 195:792–800
9. Uhlig C, Bluth T, Schwarz K, et al: Effects of volatile anesthetics on mortality and postoperative pulmonary and other complications in patients undergoing surgery: A systematic review and meta-analysis. *Anesthesiology* 2016; 124:1230–1245
10. Volta CA, Alvisi V, Petrini S, et al: The effect of volatile anesthetics on respiratory system resistance in patients with chronic obstructive pulmonary disease. *Anesth Analg* 2005; 100:348–353
11. Shamseer L, Moher D, Clarke M, et al; PRISMA-P Group: Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: Elaboration and explanation. *BMJ* 2015; 350:g7647
12. Drevon D, Fursa SR, Malcolm AL: Inter-coder reliability and validity of WebPlotDigitizer in extracting graphed data. *Behav Modif* 2017; 41:323–339
13. Wan X, Wang W, Liu J, et al: Estimating the sample mean and standard deviation from the sample size, median, range and/or interquartile range. *BMC Med Res Methodol* 2014; 14:135
14. Higgins J, Thomas J. *Cochrane handbook for systematic reviews of interventions version 6.4*. Cochrane. Published online 2023. Available at: www.training.cochrane.org/handbook. Accessed August 2023
15. Chen X, Li M, Zheng R, et al: Effects of sevoflurane inhalation anesthesia on IL-6, TNF- α and MMP-9 expression and hemodynamics in elderly patients undergoing lobectomy for lung cancer. *Cell Mol Biol (Noisy-le-grand)* 2020; 66:49–53
16. De Conno E, Steurer MP, Wittlinger M, et al: Anesthetic-induced improvement of the inflammatory response to one-lung ventilation. *Anesthesiology* 2009; 110:1316–1326
17. Delogu G, Antonucci A, Signore M, et al: Plasma levels of IL-10 and nitric oxide under two different anaesthesia regimens. *Eur J Anaesthesiol* 2005; 22:462–466
18. Erturk E, Topaloglu S, Dohman D, et al: The comparison of the effects of sevoflurane inhalation anesthesia and intravenous

- propofol anesthesia on oxidative stress in one lung ventilation. *Biomed Res Int* 2014; 2014:1–4
19. Hammouda SA, Abd Rabbih AA, AlGanady AA, et al: Immunomodulatory effect of propofol versus sevoflurane in patients undergoing thoracic surgery using one lung ventilation technique. *Egypt J Chest Dis Tuberc* 2013; 62:731–743
 20. Huang CH, Wang YP, Wu PY, et al: Propofol infusion shortens and attenuates oxidative stress during one lung ventilation. *Acta Anaesthesiol Taiwan* 2008; 46:160–165
 21. Jin Y, Zhao X, Li H, et al: Effects of sevoflurane and propofol on the inflammatory response and pulmonary function of perioperative patients with one-lung ventilation. *Exp Ther Med* 2013; 6:781–785
 22. Kim HJ, Kim E, Baek SH, et al: Sevoflurane did not show better protective effect on endothelial glycocalyx layer compared to propofol during lung resection surgery with one lung ventilation. *J Thorac Dis* 2018; 10:1468–1475
 23. Lee JJ, Kim GH, Kim JA, et al: Comparison of pulmonary morbidity using sevoflurane or propofol-remifentanyl anesthesia in an Ivor Lewis operation. *J Cardiothorac Vasc Anesth* 2012; 26:857–862
 24. Li XF, Hu JR, Wu Y, et al: Comparative effect of propofol and volatile anesthetics on postoperative pulmonary complications after lung resection surgery: A randomized clinical trial. *Anesth Analg* 2021; 133:949–957
 25. Mahmoud K, Ammar A: Immunomodulatory effects of anesthetics during thoracic surgery. *Anesthesiol Res Pract* 2011; 2011:317410
 26. Nakanuno R, Yasuda T, Hamada H, et al: Propofol for anesthesia and postoperative sedation resulted in fewer inflammatory responses than sevoflurane anesthesia and midazolam sedation after thoracoabdominal esophagectomy. *Hiroshima J Med Sci* 2015; 64:31–37
 27. Potočnik I, Novak-Janković V, Šostarič M, et al: Antiinflammatory effect of sevoflurane in open lung surgery with one-lung ventilation. *Croat Med J* 2014; 55:628–637
 28. Qiao Y, Feng H, Zhao T, et al: Postoperative cognitive dysfunction after inhalational anesthesia in elderly patients undergoing major surgery: The influence of anesthetic technique, cerebral injury and systemic inflammation. *BMC Anesthesiol* 2015; 15:154
 29. Schilling T, Kozian A, Kretzschmar M, et al: Effects of propofol and desflurane anaesthesia on the alveolar inflammatory response to one-lung ventilation. *Br J Anaesth* 2007; 99:368–375
 30. Schilling T, Kozian A, Senturk M, et al: Effects of volatile and intravenous anesthesia on the alveolar and systemic inflammatory response in thoracic surgical patients. *Anesthesiology* 2011; 115:65–74
 31. Sen Y, Xiyang H, Yu H: Effect of thoracic paraspinal block-propofol intravenous general anesthesia on VEGF and TGF- β in patients receiving radical resection of lung cancer. *Medicine (Baltim)* 2019; 98:e18088
 32. Sugawara Y, Yamaguchi K, Kumakura S, et al: Effects of sevoflurane and propofol on pulmonary inflammatory responses during lung resection. *J Anesth* 2012; 26:62–69
 33. Tian HT, Duan XH, Yang YF, et al: Effects of propofol or sevoflurane anesthesia on the perioperative inflammatory response, pulmonary function and cognitive function in patients receiving lung cancer resection. *Eur Rev Med Pharmacol Sci* 2017; 21:5515–5522
 34. Tsuchiya M, Shiimoto K, Mizutani K, et al: Reduction of oxidative stress a key for enhanced postoperative recovery with fewer complications in esophageal surgery patients: Randomized control trial to investigate therapeutic impact of anesthesia management and usefulness of simple blood test for prediction of high-risk patients. *Medicine (Baltim)* 2018; 97:e12845
 35. Von Dossow V, Baur S, Sander M, et al: Propofol increased the interleukin-6 to interleukin-10 ratio more than isoflurane after surgery in long-term alcoholic patients. *J Int Med Res* 2007; 35:395–405
 36. Wakabayashi S, Yamaguchi K, Kumakura S, et al: Effects of anesthesia with sevoflurane and propofol on the cytokine/chemokine production at the airway epithelium during esophagectomy. *Int J Mol Med* 2014; 34:137–144
 37. Wang G, Liu J, Gao J, et al: Comparison of the effects of sevoflurane and propofol anesthesia on pulmonary function, MMP-9 and postoperative cognition in patients receiving lung cancer resection. *Oncol Lett* 2019; 17:3399–3405
 38. Xu WY, Wang N, Xu HT, et al: Effects of sevoflurane and propofol on right ventricular function and pulmonary circulation in patients undergone esophagectomy. *Int J Clin Exp Pathol* 2014; 7:272–279
 39. Yang L, Wang F, Wang L: Comparison of remifentanyl with sevoflurane and remifentanyl with propofol on perioperative inflammatory response and pulmonary function in patients with lung cancer
 40. Zhang GH, Wang W: Effects of sevoflurane and propofol on the development of pneumonia after esophagectomy: A retrospective cohort study. *BMC Anesthesiol* 2017; 17:164
 41. Uchida T, Shirasawa M, Ware LB, et al: Receptor for advanced glycation end-products is a marker of type I cell injury in acute lung injury. *Am J Respir Crit Care Med* 2006; 173:1008–1015
 42. Strieter RM, Belperio JA, Keane MP: Host innate defenses in the lung: The role of cytokines. *Curr Opin Infect Dis* 2003; 16:193–198
 43. Parsons PE, Eisner MD, Thompson BT, et al: NHLBI Acute Respiratory Distress Syndrome Clinical Trials Network: Lower tidal volume ventilation and plasma cytokine markers of inflammation in patients with acute lung injury. *Crit Care Med* 2005; 33:1–6; discussion 230
 44. Martin TR: Lung cytokines and ARDS: Roger S. Mitchell Lecture. *Chest* 1999; 116(1 Suppl):2S–8S
 45. Piantadosi CA, Schwartz DA: The acute respiratory distress syndrome. *Ann Intern Med* 2004; 141:460–470
 46. Donnelly SC, Strieter RM, Reid PT, et al: The association between mortality rates and decreased concentrations of interleukin-10 and interleukin-1 receptor antagonist in the lung fluids of patients with the adult respiratory distress syndrome. *Ann Intern Med* 1996; 125:191–196
 47. Ware LB, Matthay MA: The acute respiratory distress syndrome. *N Engl J Med* 2000; 342:1334–1349
 48. Dreyfuss D, Saumon G: Ventilator-induced lung injury: Lessons from experimental studies. *Am J Respir Crit Care Med* 1998; 157:294–323

49. Schilling T, Kozian A, Huth C, et al: The pulmonary immune effects of mechanical ventilation in patients undergoing thoracic surgery. *Anesth Analg* 2005; 101:957–965
50. Jang D, Lee AH, Shin HY, et al: The role of tumor necrosis factor alpha (TNF- α) in autoimmune disease and current TNF- α inhibitors in therapeutics. *Int J Mol Sci* 2021; 22:2719
51. Costela-Ruiz VJ, Illescas-Montes R, Puerta-Puerta JM, et al: SARS-CoV-2 infection: The role of cytokines in COVID-19 disease. *Cytokine Growth Factor Rev* 2020; 54:62–75
52. Pagel PS: Myocardial protection by volatile anesthetics in patients undergoing cardiac surgery: A critical review of the laboratory and clinical evidence. *J Cardiothorac Vasc Anesth* 2013; 27:972–982
53. Pagel PS, Crystal GJ: The discovery of myocardial preconditioning using volatile anesthetics: A history and contemporary clinical perspective. *J Cardiothorac Vasc Anesth* 2018; 32:1112–1134